

A new 2012 submarine volcanic activity off the western coast of El Hierro (Canary Islands, Spain): evidences from acoustic imaging and precursory signatures

Nemesio M. Perez¹, Luis Somoza², Luis Gonzales de Vallejo⁴, Ricardo Leon², Takeshi Sagiya⁵, Pedro A. Hernandez¹, Ander Biain², Francisco J. Gonzalez², Teresa Medialdea², Daniel Gonzalez-Aller⁶, Jose L. Sanchez de La Madrid⁶, Jase Barrancos³, Jesus Ibanez⁷, Hirochika Sumino⁸, Carmen Romero⁹

¹Instituto Volcanologico de Canarias, INVOLCAN, Spain, ²Marine Geology Division, Instituto Geologico y Minero de España (IGME), Spain, ³Environmental Research Division, ITER, Spain, ⁴Facultad de Ciencias Geologicas, Universidad Complutense de Madrid, Spain, ⁵Research Center for Seismology, Volcanology, and Disaster Mitigation, Nagoya University, Japan, ⁶Instituto Hidrografico de la Marina (IHHM), Spain, ⁷Instituto Andaluz de Geofisica, Universidad de Granada, Spain, ⁸Geochemical Research Center, The University of Tokyo, Bunkyo-Ku, Japan, ⁹Facultad de Geografia e Historia, Universidad de La Laguna, Spain

E-mail: nperez@iter.es

Here we report precursory geophysical, geodetic, and geochemical signatures of a new submarine volcanic activity observed off the western coast of El Hierro, Canary Islands, which has been detected through acoustic imaging of submarine plumes on June 28, 2012, by the Spanish research vessel Hesperides. Five distinct acoustic submarine plumes have been recognized in this area at water depths between 64 and 88 m along a submarine platform located in front of the Lomo Negro volcanic cone, west of El Hierro. Submarine plumes are characterized by vertical columns of high-amplitude values rising from seafloor. These acoustic imaging data clearly support a new submarine volcanic activity in 2012 associated to the recent magmatic reactivation of El Hierro volcanic system. This new submarine volcanic activity was preceded by several precursory signatures: (i) a sharp increase of the seismic energy release and the number of daily earthquakes of magnitude higher than 2.5 on June 25, 2012, (ii) significant vertical and horizontal displacements observed at the Canary Islands GPS permanent network Nagoya University-ITER-GRAFCAN at El Hierro with uplifts up to 3 cm from June 25 to 26, 2012, (iii) an anomalous increase of the soil gas radon activity at HIE02, a geochemical station located in the northwestern of El Hierro, from the end of April until the beginning of June reaching peak values of 2.7 kBqm³ on June 3, 2012, and (iv) the highest observed corrected value of ³He/⁴He ratio in ground waters (8,5 Ra) from San Simon well at the northwestern of El Hierro on June 16, 2012. These precursory signals have revealed important to improve and optimize the detection of early warning signals of volcanic unrest episodes at El Hierro.