

## Satellite-based assessment of global volcanic degassing

Simon A. Carn<sup>1</sup>, Kai Yang<sup>2</sup>, Verity J.B. Flower<sup>1</sup>

<sup>1</sup>Dept. of Geological and Mining Eng. and Sciences, Michigan Technological University, Houghton, MI, USA, <sup>2</sup>Dept. of Atmospheric and Oceanic Sciences, University of Maryland, College Park, MD, USA

E-mail: [scarn@mtu.edu](mailto:scarn@mtu.edu)

Satellite observations by hyperspectral ultraviolet (UV) and infrared (IR) sensors over the past decade have afforded tremendous insights into the spatial and temporal variability of global, subaerial volcanic degassing. Commonly cited volcanic emissions inventories are still largely based on infrequent ground-based gas measurements and have not been updated in recent years. We use 8 years of sulfur dioxide (SO<sub>2</sub>) measurements by the Ozone Monitoring Instrument (OMI) on NASA's Aura satellite to assess the spatial and temporal variability of volcanic SO<sub>2</sub> degassing between 2004 and 2012. We focus on passive (lower tropospheric) degassing, which is the major component of total volcanic emissions to the atmosphere on a time-averaged basis, but poorly constrained. Synergy between OMI and infrared (IR) satellite sensors (e.g., Aqua/AIRS) in the A-Train satellite constellation provides constraints on the vertical distribution of SO<sub>2</sub>, which is critical for assessing the potential climate impacts of volcanic emissions. OMI measurements are most sensitive to SO<sub>2</sub> emission rates on the order of 1000 tons/day or more, and thus the satellite data provide new constraints on the location and persistence of major volcanic SO<sub>2</sub> sources. Time-averaging of OMI SO<sub>2</sub> data provides information on weaker SO<sub>2</sub> degassing. We find that OMI has detected non-eruptive SO<sub>2</sub> emissions from at least 60 volcanoes since 2004. Results of our analysis reveal the emergence of several dominant tropospheric SO<sub>2</sub> sources that are not prominent in existing inventories (Ambrym, Nyiragongo, Turrialba), the persistence of some well-known sources (Etna, Kilauea) and an apparent decline in emissions at others (e.g., Lascar). The OMI measurements provide particularly valuable information in regions lacking regular ground-based monitoring such as Indonesia, Melanesia and Kamchatka. We describe how the OMI measurements of SO<sub>2</sub> total column, and their probability density function, can be used to infer SO<sub>2</sub> emission rates for compatibility with existing datasets. We also discuss the potential biases in the OMI measurements due to latitudinal variations in ozone abundance, UV irradiance and cloud cover. Our analysis underlines the critical role of hyperspectral UV satellite observations in assessing global volcanic degassing rates, particularly for remote and/or unmonitored volcanoes.